

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

North American Crane Workshop Proceedings

North American Crane Working Group

2005

CAPTURE OF SANDHILL CRANES USING ALPHA-CHLORALOSE

MATTHEW A. HAYES

International Crane Foundation, matt@savingcranes.org

BARRY K. HARTUP

International Crane Foundation

JEANNE MARIE PITTMAN

International Crane Foundation

JEB A. BARZEN

International Crane Foundation

Follow this and additional works at: <http://digitalcommons.unl.edu/nacwgproc>



Part of the [Behavior and Ethology Commons](#), [Biodiversity Commons](#), [Ornithology Commons](#), [Population Biology Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

HAYES, MATTHEW A.; HARTUP, BARRY K.; PITTMAN, JEANNE MARIE; and BARZEN, JEB A., "CAPTURE OF SANDHILL CRANES USING ALPHA-CHLORALOSE" (2005). *North American Crane Workshop Proceedings*. 14.

<http://digitalcommons.unl.edu/nacwgproc/14>

This Article is brought to you for free and open access by the North American Crane Working Group at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in North American Crane Workshop Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

CAPTURE OF SANDHILL CRANES USING ALPHA-CHLORALOSE

MATTHEW A. HAYES,^{1,2} International Crane Foundation, E-11376 Shady Lane Road, Baraboo, WI 53913, USA

BARRY K. HARTUP, International Crane Foundation, E-11376 Shady Lane Road, Baraboo, WI 53913, USA

JEANNE MARIE PITTMAN, International Crane Foundation, E-11376 Shady Lane Road, Baraboo, WI 53913, USA

JEB A. BARZEN, International Crane Foundation, E-11376 Shady Lane Road, Baraboo, WI 53913, USA

Abstract: From 1990 – 2001, 188 captures of 166 different greater sandhill cranes (*Grus canadensis tabida*) were made through the experimental use of alpha-chloralose (AC) in Wisconsin (Hayes et al., 2003). Captures occurred in August (n = 28, 15%), September (n = 136, 72%) and October (n = 24, 13%). Capture of all members in the target social group was relatively high (59%). Territorial pairs were captured more successfully (2 of 2 cranes were captured 69% of the time) than family groups (3 of 3 cranes were captured 50% of the time, and 4 of 4 cranes were captured 59% of the time). Eighteen cranes were captured twice, and two cranes were captured three times using alpha-chloralose. There was no association between dosage provided, month of capture, or target number of cranes captured and capture success. There was also no association between sedation level the cranes experienced and the month that capture occurred. Overall morbidity (6%) and mortality (4%) were lower than most other capture techniques for sandhill cranes. Exertional myopathy was the cause of morbidity in 7 of 12 (58%) cases, and resulted in 3 deaths (all males). Rehabilitation efforts led to the recovery of 4 birds. The rehabilitated cranes were released within 2 weeks after capture, resumed normal behavior (i.e. pairs were not disrupted, etc.), and survived more than 1 year. Sedation level ($X^2_5 = 25.9$, $p < 0.01$) and month of capture ($X^2_2 = 12.3$, $p < 0.01$) were both associated with the presence of exertional myopathy in cranes captured with AC. Logistic regression analysis further suggested lighter sedation with AC and capture during the months of August and October were risk factors for exertional myopathy in this population. A decrease in sedation score by one unit increased the odds of exertional myopathy occurring by three times (i.e., the less sedated the crane, the higher the probability of developing exertional myopathy). Also, the odds of EM occurring during capture in August and October were 15 and 11 times greater compared to September, respectively. In addition to avoiding these risk factors, observing cranes after release and having the option of rehabilitation can lower mortality rates associated with AC capture of cranes.

PROCEEDINGS NORTH AMERICAN CRANE WORKSHOP 9:248

Key words: capture techniques, exertional myopathy, *Grus canadensis*, morbidity, mortality, sandhill crane

LITERATURE CITED

- Hayes, M. A., B. K. Hartup, J. M. Pittman, J. A. Barzen. 2003.
Capture of sandhill cranes using alpha-chloralose. *Journal of Wildlife Diseases* 39:859-868.

¹ Current address: University of South Dakota, Department of Biology, 414 East Clark Street, Vermillion, SD 57069, USA

² Corresponding author, e-mail: matt@savingcranes.org